Cefas

Cefas Environment Report RL 14/2009

Radiological assessment of dredging application for Oldbury power station (2009)

Cefas Contract SLBA1 Marine material disposal – Part II FEPA

Environment Report RL 14 /2009

RADIOLOGICAL ASSESSMENT OF DREDGING APPLICATION FOR OLDBURY POWER STATION (2009)

The Centre for Environment, Fisheries & Aquaculture Science
Lowestoft Laboratory
Pakefield Road
Lowestoft
Suffolk
NR33 0HT

K.S. Leonard

March 2009

The work in this report was completed under contract Defra

Cefas Contract SLBA1 Marine material disposal – Part II FEPA

RADIOLOGICAL ASSESSMENT OF DREDGING APPLICATION FOR OLDBURY POWER STATION (2009)

Summary

In 2009, Magnox North Limited lodged a FEPA licensing application to carry out a 3 months dredging program, involving the disposal at sea of 360 m^3 of sediment from Oldbury Power Station. Using the conservative generic radiological assessment procedure developed by the IAEA, to convert radionuclide concentrations in dumped material into radiation doses due to dumping, the total derived total dose to individual members of the crew and public was 3.0 µSv/year and ~1.0 µSv/year, respectively. The total collective dose was 0.026 manSv/year. The values for individual members of the crew and public, and the collective dose, were within the *de minimis* criteria of 10 µSv/year (individual doses) and 1 manSv/year (collective dose), respectively.

Since the conservative generic radiological assessment procedure indicated that doses received were well below recommended limits, a subsequent more detailed case specific assessment was not necessary. Therefore, from radiological considerations, there is no objection to this material being dredged and dumped.

Assessment details

In 2009, Magnox North Limited lodged a FEPA licensing application to carry out a dredging program (3 months) involving the disposal at sea of 360 m^3 of sediment from Oldbury Power Station. As part of the radiological assessment of the proposed operation, four sediment samples (S1 –S4, inclusive) were collected in February 2009. Sampling locations were:

S1 - North Dredge Area A, S2 - North Dredge Area B, S3 - South Dredge Area A, and S4 - South Dredge Area B.

These locations are illustrated in Figure 1, together with the proposed offshore disposal "box" area.

Figure 1. Study Area (Oldbury Power Station) and Sampling Locations (S1 –S4)



Following freeze-drying and homogenisation, radionuclide assay at the Cefas Lowestoft Laboratory was achieved by γ counting samples on a high purity Ge detector. The results are summarised in Table 1.

Table 1. Radioactivity in sediment dredged from Oldbury Power Station, February 2009

Sample Location	Specific activity (Bq/kg, dry weight)					
	^{60}Co	^{137}Cs	^{226}Ra (via ^{214}Pb)	^{232}Th (via ^{228}Ac)	^{238}U (via ^{234}Th)	^{241}Am
S1	< 0.25	12	17	19	30	< 1.1
S2	< 0.28	8	13	15	19	< 0.45
S3	< 0.27	15	24	26	37	< 1.24
S4	< 0.40	6	15	16	19	< 0.68
***Average**	**0.3**	**10**	**17**	**19**	**26**	**1.0**

*Average determinations use < results as positively measured values to produce a conservative estimate

The specific activity of the artificial radionuclides (e.g. ^{137}Cs and ^{241}Am) in these samples was typical of muddy sediments from the combined effects of discharges from the site, other nuclear establishments discharging into the Bristol Channel and weapons testing (and possibly a small Sellafield derived component), being enhanced above background levels outside the Irish Sea. The Sellafield component is a legacy of large discharges from the Sellafield Limited reprocessing plant (formally British Nuclear Fuels) at Sellafield in the 1970s.

In addition to the nuclides detected by gamma spectrometry, sediment in Oldbury is known to contain activities of Pu radionuclides. The ^{241}Am data were used to derive estimates for ^{239,240}Pu and ^{241}Pu, assuming their activity was proportional to the ratio in the time integrated Sellafield discharges. This approach is reasonable given that both radionuclides are highly particle-reactive, hence the fate following discharge is similar. The activity for ^{210}Pb was derived using data for ^{226}Ra and assuming secular equilibrium.

Under the London Convention, only materials with *de minimis* levels of radioactivity may be considered for dumping. Using the conservative generic radiological assessment procedure developed by the IAEA (IAEA, 2004), to convert radionuclide concentrations in dumped material into radiation doses due to dumping, the total derived total dose to individual members of the crew and public was 3.0 µSv/year and 0.8 µSv/year, respectively. The total collective dose was 0.026 manSv/year. The values for individual members of the crew and public, and the collective dose, were within the *de minimis* criteria of 10 µSv/year (individual doses) and 1 manSv/year (collective dose), respectively.

The dose estimates for individual crew/public (by nuclide), derived using the generic IAEA model, are shown in Figure 2.

Figure 2. Assessment of dose to individual members of crew and the public arising from at Oldbury Power station a) Doses were derived using average activities listed in Table 1.



Since the conservative generic radiological assessment procedure indicated that doses received were well below recommended limits, a subsequent more detailed case specific assessment was not necessary. All the derived total dose values were less than the *de minimis* criteria of 10 µSv/a and 1 manSv/a for individual and collective dose, respectively.

Therefore, from radiological considerations, there is no objection to this material being dredged and dumped.

References

IAEA(2004). Sediment distribution coefficients and concentration factors for biota in the marine environment. Tech. Rep. Ser. No. 422, IAEA, Vienna.

About us

Cefas is a multi-disciplinary scientific research and consultancy centre providing a comprehensive range of services in fisheries management, environmental monitoring and assessment, and aquaculture to a large number of clients worldwide.

We have more than 500 staff based in 2 laboratories, our own ocean-going research vessel, and over 100 years of fisheries experience.

We have a long and successful track record in delivering high-quality services to clients in a confidential and impartial manner.
(www.cefas.co.uk)

Cefas Technology Limited (CTL) is a wholly owned subsidiary of Cefas specialising in the application of Cefas technology to specific customer needs in a cost-effective and focussed manner.

CTL systems and services are developed by teams that are experienced in fisheries, environmental management and aquaculture, and in working closely with clients to ensure that their needs are fully met.
(www.cefastechnology.co.uk)

Customer focus

With our unique facilities and our breadth of expertise in environmental and fisheries management, we can rapidly put together a multi-disciplinary team of experienced specialists, fully supported by our comprehensive in-house resources.

Our existing customers are drawn from a broad spectrum with wide ranging interests. Clients include:

- international and UK government departments
- the European Commission
- the World Bank
- Food and Agriculture Organisation of the United Nations (FAO)
- oil, water, chemical, pharmaceutical, agro-chemical, aggregate and marine industries
- non-governmental and environmental organisations
- regulators and enforcement agencies
- local authorities and other public bodies

We also work successfully in partnership with other organisations, operate in international consortia and have several joint ventures commercialising our intellectual property.

Head office
Centre for Environment,
Fisheries & Aquaculture Science
Pakefield Road, Lowestoft,
Suffolk NR33 0HT UK

Tel +44 (0) 1502 562244
Fax +44 (0) 1502 513865
Web www.cefas.co.uk

printed on paper made from a minimum 75% de-inked post-consumer waste